

### Control of Weeds: grasses & broad leaf



#### Rax Super 7.5EW

- Outstanding and economic management of a wide range of annual grasses.
- Contact and systemic herbicide.
- Flexibility in application timing.
- Excellent rain fastness properties
- More stable formulation.

Rate: 0.8-1L/Ha, 100mls/20L  
Pack Sizes: 1L, 5L



### Control Of Broad Leaf Weeds In Wheat And Barley



#### Ariane

- It is also highly efficient in the selective elimination.
- control of an expansive range of weeds.
- It is versatile in its use for both high altitude and low altitude broad-leaved weed control.

Rates: 80-100mls/20L (1.25L/Ha)  
Pack Size: 1L, 5L



### Control of Sucking Pests (Russian Aphids & Rose Grain Aphids)



#### Blast 60EC

- Broad spectrum of activity for the control of soil borne insects (Wireworms) and sucking insects e.g. Russian aphids.
- Quick knockdown effect.
- Easily absorbed in plant tissues and digested by the target pest.
- Long lasting residual control.

NB: 1st spray to control soil-borne insects (wireworms). The product may be tank-mixed with most used herbicide.

It is important to rotate the pesticides to avoid pest resistance.

Rate: 0.75L/Ha, 30mls/20L  
Pack Sizes: 500ml, 1L



### Control Of Sucking Pest: Aphids & Spider Mites



#### Cutter 112EC

- Broad-spectrum insecticide to control sucking insects.
- Low application rates making it economical.
- Contact and systemic action.
- Quick knock down effect.
- Long lasting residual action.

Rates: 350mls/Ha, 20mls/20L  
Pack Sizes: 100ml, 500ml



### Control of Rust (Stem, Yellow and Leaf)



#### Culture Control Methods:

- All the three rusts are primarily wind-borne; they are not seed or soil-borne.
- Avoid planting very susceptible wheat varieties - Use resistant varieties wherever possible.
- Remove or destroy volunteer wheat plants after the season.
- Monitor the crop for signs of infection from 60 days or two months after planting.
- Early fungicide application/protection is important.
- Crop rotation with crops like canola and beans.

#### April 250 EC

- Broad-spectrum fungicide for the control of leaf, stem and yellow rust.
- Highly effective Tebuconazole in the market.

Rate: 0.75 L/Ha, 75mls/20L  
Pack Sizes: 1L, 5L



#### Othello Top

- Preventive and Curative function.
- Long lasting residual protection.
- Highly systemic.
- Excellent rain fast ability due to special emulsifiers.

Rates: 0.5L/Ha, 50mls/20L  
Pack Sizes: 1L, 5L



### Control of Fusarium Head Blight (FHB)



#### Culture Control Methods:

- Symptoms are confined to the head, grain and sometimes on the peduncle (neck).
- Plant resistant varieties.
- Crop rotation, e.g. Canola or Potatoes.
- Apply a foliar fungicide (Orius) early to prevent infection at flowering.

#### April 250 EC

- Broad-spectrum fungicide for the control of leaf, stem and yellow rust.
- The highly rated effective Tebuconazole in the market.
- Cost effective. Preventive and curative action against a wide range of leaf diseases.

Rate: 0.75 L/Ha, 75mls/20L  
Pack Sizes: 1L, 5L

